

TWO-PARAMETRIC QUANTUM-DEFORMED DUAL AMPLITUDE

L.Jenkovszky

Bogoliubov Institute for Theoretical Physics
Metrologicheskaya St. 14b, 252143, Kiev-143, Ukraine.
E-mail:jenk@gluk.apc.org

M.Kibler

Institut de Physique Nucléaire de Lyon
IN2P3-CNRS et Université Claude Bernard
43 Boulevard du 11 Novembre 1918
F69622 Villeurbanne Cedex, France
E-mail:kibler@frcpn11.bitnet

A.Mishchenko

Bogoliubov Institute for Theoretical Physics
Metrologicheskaya St. 14b, 252143, Kiev-143, Ukraine.
E-mail:yusitenko@gluk.apc.org

February 1, 2008

Abstract

From a 2-parametric deformation of the harmonic oscillator algebra we construct a 4-point dual amplitude with nonlinear trajectories. The earlier versions of the q-deformed dual models are reproduced as limiting cases of the present model.

Originally, dual models were thought to be promising examples of hadronic scattering amplitudes in the framework of the S -matrix theory. Subsequently, they gave rise to many formal and mathematical developments while the original goal of constructing realistic scattering amplitudes was abandoned. One of the difficult problems that remained unsolved is that of the spectrum of particles in the model and the related problem of the form of the Regge trajectories used.

We remind that in the original Veneziano amplitude [1] the trajectories are linear and the model predicts the existence of an infinite sequence of infinitely narrow "resonances" whose spin increases like their squared masses. Narrow resonance dual models have the virtue of being factorizable [2], giving rise to a field-theoretical interpretation of the scattering amplitude in terms of relevant vertices and propagators, which in principle can be used in the construction of a unitarization procedure (calculation of loop diagrams) based on the Veneziano dual amplitude as the "Born term". In spite of many efforts with partial successes, this program has never been completed.

A different class of dual models [3-5] is based on logarithmic trajectories. The existence of ghosts - states with negative norm - was one of the main reason why the study of this second class of dual models was abandoned.

Factorizability made it possible for both classes of the dual models to construct an operator formalism corresponding to an infinite set of oscillators. It has been also realized that the algebra of the dual models with logarithmic trajectories can be viewed as the q -deformed algebra of harmonic oscillators (Veneziano model).

Recent interest in quantum algebras (q -deformed algebras) brought also a revival to the traditional dual models. Various q -deformations of the harmonic oscillators and the resulting "dual amplitudes" have been studied in a number of papers [4,6,7].

In the present paper we construct a quantum-deformed dual amplitude by means of a 2-parametric deformation of the dual amplitude.

We remind that a similar problem in the case of a single-parameter deformation was considered in refs. [4,6,7]. The aim of all these attempts was to replace the infinite-dimensional algebra of the independent harmonic oscillators by a q -deformed quon algebra [8] which essentially means the introduction of some anharmonism in the interaction [9]. Correspondingly are the vertices and the propagators also modified.

In refs. [4,7], the following deformation was used:

$$a_{\mu}^{(n)} a_{\nu}^{+(n)} - q a_{\nu}^{+(n)} a_{\mu}^{(n)} = -\eta_{\mu\nu}, \quad n = \overline{1, \infty}, \quad (1)$$

where $\eta_{\mu\nu} = \text{diag}(1, -1, \dots, -1)$, $|q| < 1$.

In ref. [6] on the other hand, the q deformation was defined as

$$\begin{aligned} a_{\mu}^{(n)} a_{\nu}^{+(n)} - q a_{\nu}^{+(n)} a_{\mu}^{(n)} &= -\eta_{\mu\nu} q^{-N_n}, \\ [N_n, a_{\mu}^{(n)}] &= -a_{\mu}^{(n)}, \quad [N_n, a_{\mu}^{+(n)}] = a_{\mu}^{+(n)}. \end{aligned} \quad (2)$$

Note that in eqs. (1) and (2) q is a real parameter and the interaction between various modes n and m is absent:

$$[a_{\mu}^{(n)}, a_{\mu}^{(m)}] = [a_{\mu}^{(n)}, a_{\mu}^{+(m)}] = [a_{\mu}^{+(n)}, a_{\mu}^{+(m)}] = 0, \quad n \neq m. \quad (3)$$

The 2-parametric deformation to be introduced in the present paper generalizes these two approaches and contains them as limiting cases.

Let us remind that the 4-point dual Veneziano amplitude is defined by the following expression

$$A_4(s, t) = B(-\alpha_0(s), -\alpha_0(t)) = \int_0^1 dx x^{-\alpha_0(s)-1} (1-x)^{-\alpha_0(t)-1} \\ = \langle 0 | V(p_2) \int_0^1 dx x^{H-\alpha_0(s)-1} (1-x)^{\alpha(0)-1} V(p_3) | 0 \rangle, \quad (4)$$

where α_0 is a (linear) Regge trajectory

$$\alpha_0(s) = \alpha' s + \alpha(0), \quad p_i^2 = m^2, \quad \alpha' m^2 + \alpha(0) = 0, \\ 2\alpha'(p_2 p_3) = \alpha' t + 2\alpha(0), \\ 2\alpha'(p_1 p_2) = \alpha' s + 2\alpha(0), \quad (5)$$

depending on the invariant (Mandelstam) kinematical variables

$$s = (p_1 + p_2)^2, \quad t = (p_2 + p_3)^2, \quad u = (p_1 + p_3)^2.$$

The relevant vertex operator and the Hamiltonian are

$$V(p) = \exp(-\sqrt{2\alpha'} p^\mu \sum_{n=1}^{\infty} \frac{a_\mu^{+(n)}}{\sqrt{n}}) \exp(\sqrt{2\alpha'} p^\mu \sum_{n=1}^{\infty} \frac{a_\mu^{(n)}}{\sqrt{n}}), \\ H = - \sum_{n=0}^{\infty} n a_\mu^{+(n)} a^{\mu(n)} \quad (6)$$

and the creation and annihilation operators a_μ^n , $a_\mu^{+(n)}$ in (6) satisfy the commutation relations (1) for $q=1$.

For $\alpha(0) = 1$, the amplitude (4) exhibits remarkable factorization properties, as one can easily verify by integrating (4) over x :

$$A_4(s, t) = \langle 0 | V(p_2) [H - \alpha_0(s)]^{-1} V(p_3) | 0 \rangle. \quad (7)$$

From (7) the pole structure of the amplitude can be easily found; the poles are located at

$$\alpha_0(s_n) = \alpha' s_n + \alpha(0) = E_n, \quad s_n = \frac{E_n - \alpha(0)}{\alpha'}, \quad (8)$$

where E_n is the eigenvalue of the operator H .

Crossing symmetry and duality imply [10,11]

$$A_4(s, t) = - \frac{(\alpha_0(s) + \alpha_0(t))}{\alpha_0(s)\alpha_0(t)} \prod_{n=1}^{\infty} \frac{n[n - \alpha_0(s) - \alpha_0(t)]}{[n - \alpha_0(s)][n - \alpha_0(t)]} \\ = - \sum_{n=0}^{\infty} \frac{(\alpha_0(s) + 1) \dots (\alpha_0(s) + n)}{n!} \frac{1}{\alpha_0(t) - n} \\ = A_4(t, s) = - \sum_{n=0}^{\infty} \frac{(\alpha_0(t) + 1) \dots (\alpha_0(t) + n)}{n!} \frac{1}{\alpha_0(s) - n}. \quad (9)$$

In deforming the Veneziano amplitude $A_4(s, t)$, we use the following quon algebra [12]:

$$\begin{aligned} a_\mu^{(n)} a_\nu^{+(n)} - q_1 a_\nu^{+(n)} a_\mu^{(n)} &= -\eta_{\mu\nu} q_2^{-N_n}, \quad \mu, \nu = \overline{1, d}, \quad n = \overline{1, \infty} \\ [N_n, a_\mu^{(n)}] &= -a_\mu^{(n)}, \quad [N_n, a_\mu^{+(n)}] = a_\mu^{+(n)}, \\ [a_\mu^{(n)}, a_\nu^{(m)}] &= [a_\mu^{(n)}, a_\nu^{+(m)}] = [a_\mu^{+(n)}, a_\nu^{+(m)}] = 0, \quad m \neq n, \end{aligned} \quad (10)$$

where q_1 and q_2 are real numbers whose moduli do not belong to $[0, 1]$ simultaneously.

Introduce now the following notations, usefull in what follows

$$\begin{aligned} a^{(n)}(p) &= a_\mu^{(n)} p^\mu, \quad a^{+(n)}(p) = a_\mu^{+(n)} p^\mu, \\ H_n &= a_\mu^{+(n)} a^{\mu(n)}, \end{aligned} \quad (11)$$

and

$$(x)_q = \frac{q_1^x - q_2^x}{q_1 - q_2}, \quad (12)$$

where p^μ is a c-number vector and x is a number or operator.

To derive the deformed amplitude, first we need the construction of the representation of the quon algebra (10). The precEDURE is standard and is based on the introduction of the vacuum (cyclic vector $|0\rangle$):

$$a_\mu^{(n)} |0\rangle = 0, \quad N |0\rangle = 0, \quad (13)$$

as well as the state vectors - by applying the creation operators $a_\mu^{+(n)}$ on the vacuum.

We are interested only in Lorentz-invariant combinations (11), for which the relations

$$\begin{aligned} a^{(n)}(p_1) a^{+(n)}(p_2) - q_1 a^{+(n)}(p_2) a^{(n)}(p_1) &= -(p_1 p_2) q_2^{N_n}, \\ H_n a^{+(n)}(p) |0\rangle &= -a^{+(n)}(p) |0\rangle. \end{aligned} \quad (14)$$

hold. By induction, the following relations can be easily derived from (10) and (14):

$$\begin{aligned} H_n [a^{+(n)}(p)]^k |0\rangle &= -(k)_q [a^{+(n)}(p)]^k |0\rangle \\ \langle 0 | [a^{(n)}(p_1)]^k [a^{+(n)}(p_2)]^l |0\rangle &= \delta_{kl} (-1)^k [(k)_q!] (p_1 p_2)^k, \end{aligned} \quad (15)$$

where

$$(0)_q! = 1, \quad (k)_q! = \prod_{j=1}^k (j)_q!, \quad k \geq 1. \quad (16)$$

The relations (15) will be basic when calculating the deformed amplitude.

Our objective is the construction of a deformed dual and factorizable scattering amplitude. Any factorizable representation for the scattering amplitude will require the calculation of the vacuum expectation values of the form

$$\begin{aligned}
& < 0 | V(a^{(n)}(p_1)) D(H_n) V(-a^{+(n)}(p_2)) | 0 > \\
& = \sum_{n=0}^{\infty} V_1^{(n)} V_2^{(n)} [(n)_q!] (p_1 p_2)^n D(-(n)_q),
\end{aligned} \tag{17}$$

where

$$V_i(x) = \sum_{n=0}^{\infty} V_i^{(n)} x^n. \tag{18}$$

Note that eq. (17) is invariant under the replacement $H_n \longrightarrow -(N_n)_q$.

Now, following [6,7] we define a new vertex operator replacing the usual exponential by a deformed one

$$e_q(x) = \sum_{n=0}^{\infty} \frac{x^n}{[(n)_q!]}, \tag{19}$$

satisfying the equation

$$\frac{e_q(q_1 x) - e_q(q_2 x)}{x(q_1 - q_2)} = e_q(x). \tag{20}$$

It is used in the constructin of the coherent states for the deformed oscillator algebra [13]

$$aa^+ - q_1 a^+ a = q_2^{-N}. \tag{21}$$

Really, it can be easily shown with the help of (20) and (21) that

$$a | z > = z | z >, \quad | z > = e_q(za^+) | 0 >. \tag{22}$$

The new vertex is now

$$V(p) = \prod_{n=1}^{\infty} e_q\left(-\frac{\sqrt{2\alpha'} a^{+(n)}(p)}{\sqrt{n}}\right) e_q\left(\frac{\sqrt{2\alpha'} a^{(n)}(p)}{\sqrt{n}}\right), \tag{23}$$

It reduces to a non-deformed one as $q_1 \rightarrow 1$ and $q_2 \rightarrow 1$.

Now by analogy with (6), we define

$$H = - \sum_{n=0}^{\infty} n a_{\mu}^{+(n)} a^{\mu(n)} = - \sum_{n=0}^{\infty} n H_n, \tag{24}$$

and by using (17) we find

$$\begin{aligned}
& < 0 | V(p_2) x^H V(p_3) | 0 > = F(x, p_2 p_3) \\
& = \prod_{n=1}^{\infty} \left(\sum_{m=0}^{\infty} \frac{1}{(m)_q!} \left(\frac{2\alpha'}{n} (p_2 p_3) \right)^m x^{n(m)_q} \right).
\end{aligned} \tag{25}$$

It can be easily verified that the insertion of (25) into (4) violates crossing symmetry. To restore symmetry, we can either define the propagator as [7]

$$D(p_1 p_2) = \int_0^1 dx x^H F(1-x, p_1 p_2), \tag{26}$$

and

$$A_4 = < 0 | V(p_2) D(p_1 p_2) V(p_3) | 0 >, \tag{27}$$

or, following ref. [6] define a set of the operators $b_\mu^{(n)}, b_\mu^{+(n)}$ commuting with $a_\mu^{(n)}, a_\mu^{+(n)}$ and similar to $a_\mu^{(n)}, a_\mu^{+(n)}$ in the sense of the mutual commutation relations (i.e. satisfying (10) under the replacement $a_\mu^{(n)} \longrightarrow b_\mu^{(n)}$ and $a_\mu^{+(n)} \longrightarrow b_\mu^{+(n)}$).

The next step towards the restoration of duality and crossing symmetry is to use the representation [6]

$$A_4 = \int_0^1 dx < 0 | V_a(p_2) V_b(p_1) x^{H_a} (1-x)^{H_b} V_b(p_2) V_a(p_3) | 0 >, \quad (28)$$

where the indices a and b mean that in the corresponding function the arguments are operators $a_\mu^{(n)}, a_\mu^{+(n)}$ or $b_\mu^{(n)}, b_\mu^{+(n)}$. The two approaches result in the same result for the deformed amplitude

$$A_4(p_2 p_3, p_1 p_2) = \int_0^1 dx F(x, p_2 p_3) F(1-x, p_1 p_2), \quad (29)$$

It can be seen from (25) that eq. (29) reduces to a non-deformed amplitude (4) for $\alpha(0) = 1$ in the limit $q_1 \rightarrow 1$ and $q_2 \rightarrow 1$, to the one given in [6] when $q_1 = q$, $q_2 = q^{-1}$ and that of ref. [7] when $q_1 = q$, $q_2 = 1$.

Let us now find the position of the poles and the Regge trajectory for the amplitude (29). To this end one has to study the behaviour of the integrand in (29) and consequently - the convergence of the infinite product (25) defining the function $F(x, p_2 p_3)$. Rewrite (25) in the form

$$F(x, p_2 p_3) = \prod_{n=1}^{\infty} (1 + c_n(x)),$$

$$c_n(x) = \sum_{m=1}^{\infty} \frac{1}{(m)_q!} \left(\frac{2\alpha'}{n} (p_2 p_3) \right)^m x^{n(m)_q}, \quad (30)$$

suitable for the utilisation of the familiar convergence criterion of infinite products [11], according to which a product of the type (30) converges absolutely if the series

$$\sum_{n=1}^{\infty} c_n(x) \quad (31)$$

converges absolutely.

To study the convergence of the series (31) with the coefficients $c_n(x)$, defined by (30), we first estimate the ratio

$$\begin{aligned} \frac{|c_{n+1}(x)|}{|c_n(x)|} &= \frac{\sum_{m=1}^{\infty} \frac{1}{(m)_q!} \left| \frac{2\alpha'}{n+1} (p_2 p_3) \right|^m x^{(n+1)(m)_q}}{\sum_{m=1}^{\infty} \frac{1}{(m)_q!} \left| \frac{2\alpha'}{n} (p_2 p_3) \right|^m x^{n(m)_q}} \\ &\leq |x| \frac{\sum_{m=1}^{\infty} \frac{1}{(m)_q!} \left| \frac{2\alpha'}{n+1} (p_2 p_3) \right|^m x^{n(m)_q}}{\sum_{m=1}^{\infty} \frac{1}{(m)_q!} \left| \frac{2\alpha'}{n} (p_2 p_3) \right|^m x^{n(m)_q}} \leq |x|. \end{aligned} \quad (32)$$

It follows from (32) that the series (31), and consequently the infinite product (30) converge absolutely for $x \in [0, 1]$. Hence, the integrand in (29) may have singularities only at the endpoints of the integration $x = 0$ and $x = 1$.

To determine the behaviour of the function $F(x, p_2 p_3)$ at $x \rightarrow (1 - 0)$, we write for the first term in the expansion in powers of $(p_2 p_3)$:

$$\ln F(x, p_2 p_3) = -2\alpha'(p_2 p_3) \ln(1 - x) + \sum_{k=2}^{\infty} [2\alpha'(p_2 p_3)]^k F_k(x). \quad (33)$$

From (33) it follows that

$$F(x, p_2 p_3) \sim (1 - x)^{-2\alpha'(p_2 p_3)}, \quad x \rightarrow (1 - 0). \quad (34)$$

Now, by using (25), (29) and (34), we find that the poles of the amplitude (29) in $p_2 p_3$ will be located at

$$2\alpha'(p_2 p_3) = n_0 + 1 + \sum_{i=1}^{\infty} i(n_i)_q, \quad n_0, n_i \in Z^+, \quad Z^+ = 0 \cup N, \quad (35)$$

where N is the set of natural numbers. Here the term n_0 comes from the expansion of

$$F(x, p_2 p_3)(1 - x)^{2\alpha'(p_2 p_3)},$$

in powers of $(1 - x)$, the term 1 comes from the integration over x in (29) and the last term in (35) comes from $F(1 - x, p_1 p_2)$.

The residue at the pole (35) is a polynomial of power

$$\sum_{i=1}^{\infty} n_i, \quad (36)$$

in $(p_1 p_2)$ (with n_i from (35)). Then we find from (35) and (36) the leading Regge trajectory in an implicit form

$$(J)_q = \frac{q_1^J - q_2^J}{q_1 - q_2} = m^2 - 1, \quad (37)$$

or (see also [14])

$$(\alpha(t))_q = \frac{q_1^{\alpha(t)} - q_2^{\alpha(t)}}{q_1 - q_2} = \alpha' t + 2\alpha(0) - 1. \quad (38)$$

In the case $q_1 = q$, $q_2 = q^{-1}$ we obtain from (37), (38) the following spin-squared mass relation

$$J(m^2) = \frac{1}{\ln q} \ln \frac{[(q - q^{-1})(m^2 - 1) + \sqrt{(q - q^{-1})^2(m^2 - 1)^2 + 4}]}{2}, \quad (39)$$

or Regge trajectory

$$\alpha(t) = \frac{1}{\ln q} \ln \frac{[(q - q^{-1})(\alpha' t + 2\alpha(0) - 1) + \sqrt{(q - q^{-1})^2(\alpha' t + 2\alpha(0) - 1)^2 + 4}]}{2}, \quad (40)$$

while for $q_1 = q$, $q_2 = 1$ we get a logarithmic trajectory

$$J(m^2) = \frac{\ln[(q - 1)(m^2 - 1) + 1]}{\ln q}, \quad (41)$$

or

$$J(m^2) = \frac{\ln[(q - 1)(\alpha' t + 2\alpha(0) - 1) + 1]}{\ln q}, \quad (42)$$

in agreement with [4].

The algebra (10) is not very convenient since nothing can be said about the commutators of the type $[a_\mu^{(n)}, a_\nu^{(n)}]$ and $[a_\mu^{+(n)}, a_\nu^{+(n)}]$. For this reason, it is of interest to find a deformed algebra of the oscillators $d_\mu^{(n)}, d_\mu^{+(n)}$ resulting in the same amplitude for the substitution $a_\mu^{(n)} \rightarrow d_\mu^{(n)}, a_\mu^{+(n)} \rightarrow d_\mu^{+(n)}$ but with known comutators between the elements of the algebra. It can be easily varified that the relevant algebra is

$$\begin{aligned} d_\mu^{(n)} d_\nu^{+(n)} - \left[\frac{(N_n + 1)_q}{(N_n)_q} \frac{N_n}{N_n + 1} \right] d_\nu^{+(n)} d_\mu^{(n)} &= -\eta_{\mu\nu} \frac{(N_n + 1)_q}{N_n + 1}, \\ [d_\mu^{(n)}, d_\nu^{(n)}] &= [d_\mu^{+(n)}, d_\nu^{+(n)}] = 0, \\ [N_n, d_\mu^{(n)}] &= -d_\mu^{(n)}, [N_n, d_\mu^{+(n)}] = d_\mu^{+(n)}, \quad n = \overline{1, \infty}, \end{aligned} \quad (43)$$

(the operators corresponing to different modes $n \neq m$ commute) since eq. (17) is invariant under the replacement $a_\mu^{(n)} \rightarrow d_\mu^{(n)}, a_\mu^{+(n)} \rightarrow d_\mu^{+(n)}$.

Algebra (43) admits for a representation by means of the convential (non-deformed) oscillators

$$\begin{aligned} d_\mu^{(n)} &= \sqrt{\frac{(N_n + 1)_q}{N_n + 1}} \tilde{d}_\mu^{(n)}, \\ d_\mu^{+(n)} &= \tilde{d}_\mu^{+(n)} \sqrt{\frac{(N_n + 1)_q}{N_n + 1}}, \end{aligned} \quad (44)$$

where

$$\begin{aligned} [\tilde{d}_\mu^{(n)}, \tilde{d}_\nu^{+(n)}] &= -\eta_{\mu\nu}, \\ [\tilde{d}_\mu^{(n)}, \tilde{d}_\nu^{(n)}] &= [\tilde{d}_\mu^{+(n)}, \tilde{d}_\nu^{+(n)}] = 0, \\ N_n &= \tilde{d}_\mu^{+(n)} \tilde{d}_\mu^{(n)}, \end{aligned} \quad (45)$$

and the opetors corresponding to various modes $n \neq m$ commuting among each other.

Interestingly, algebra (43) is a special case of the deformed algebra

$$\begin{aligned} d_\mu^{(n)} d_\nu^{+(n)} - \varphi(N_n) d_\nu^{+(n)} d_\mu^{(n)} &= -\eta_{\mu\nu} g(N_n), \\ [d_\mu^{(n)}, d_\nu^{(n)}] &= [d_\mu^{+(n)}, d_\nu^{+(n)}] = 0, \\ [d_\mu^{(n)}, d_\nu^{+(m)}] &= 0, \quad n \neq m, \end{aligned} \quad (46)$$

the latter being a generalization of the deformed algebra suggested recently [15] for the one-dimensional case.

Recently a very interesting possibility for the solution of the problem of spectrum in dual models has been suggested [16]. These authors use the general definition of duality (9) without applying to the technique of q -deformations.

In conclusion we note that all the attempts to deform the Veneziano model (harmonic oscillator) imply tree diagrams, or Born amplitudes, to be unitarized (see the beginning of this paper). In an alternative approach to the dual theory called dual amplitudes with Mandelstam analyticity [17], the input scattering amplitude already has analytic properties expected from planar unitarization (i.e. broad resonances, Mandelstam analyticity etc.).

To find the relevant operator formalism (deformed algebra?) is a challenge for the theory.

We thank A.Bugrij and B.Struminsky for useful discussions.

The work of A.M. was supported in part by a Soros Foundation grant awarded by the American Physical Society and by International Science Foundation grant.

REFERENCES

1. G.Veneziano, Nuovo Cimento, **57A** (1968) 190.
2. S.Fubini, D.Gordon and G.Veneziano, Phys. Letters, **29B** (1969) 679.
3. D.D.Coon and M.Baker, Phys. Rev., **D2** (1970) 2349.
4. D.Coon, S.Yu and M.Baker. Phys.Rev. **D5** (1972) 1429.
5. E.Gremmer and J.Nuyts, Nuc. Phys., **B26** (1971) 151.
 J.M.Golden, Lett. Nuovo Cim., (Ser.2) **1** (1971) 893.
 S.Machida, Progr. Theor. Phys., **47** (1972) 2015.
 L.J.Romans, in Proceedings of the High Energy Physics Cosmology Conference, Trieste, (1989), eds. J.C.Pati et al., World Scientific (1990).
6. M.Chiachian, J.F.Gomes and P.Kulish, Phys. Letters, **B311** (1993) 93.
7. L.L.Jenkowszky, A.V.Mishchenko, B.V.Struminsky. Preprint ITP-93-36E and in Proc. of the HADRONS-93.
8. O.W.Greenberg, Phys. Rev. Lett., **64** (1990) 705; Phys. Rev., **D43** (1991) 4111.
 R.N.Mohapatra, Phys. Lett., **B242** (1990) 407.
9. S.V.Shabanov, Phys. Lett., **B293** (1992) 117; J. Phys., **A26** (1993) 2583.
10. M.B.Green, J.H.Schwarz and E.Witten, Superstring Theory (Cambridge Univ. Press, Cambridge, 1987).
11. E.T.Whittaker, G.N.Watson, A Course of Modern Analysis (Cambridge, At the University Press, 1927).
12. M.Kibler, in Symmetry and Structural Properties of Condensed Matter, ed. W.Florek, D.Lipinski and T.Lulek, World Scientific: Sigapore (1993) 445; Miscellaneous Physical Applications of Quantum Algebras, Preprint LYCEN/9358.
13. D.Ellinas, J. Phys., **A26** (1993) L543; On Coherent States and q-Deformed Algebras, Preprint FTUV/93-37 and IFIC/93-20.
14. M.Chiachian, J.F.Gomes and R.Gonzalez Felipe, New Phenomenon of Nonlinear Regge Trajectory and Quantum Dual String Theory, Preprint HU-SEFT R 1994-05.
15. S.Meljanac, M.Mileković and S.Pallua, Unified View of Deformed Single-Mode Oscillator Algebras, Preprint RBI-TH-3/94 and PMF-ZTF-3/94 (to appear in Phys. Lett **B**).
16. D.B.Fairlie, J.Nuyts, A Fresh Look at Generalized Veneziano Amplitudes, Preprint DTP/94/19.
17. A.I.Bugrij, G.Cohen-Tannoudji, L.L Jenkowszky, N.A.Kobylinsky, Fortschritte Phys., **21** (1973) 427.